

Spectral Evolution of the Peculiar Ic Supernova 1998bw

R. A. Stathakis,¹ B. J. Boyle,¹ D. H. Jones,² M. S. Bessell,² T. J. Galama,³ Lisa M. German,⁴

¹*Anglo-Australian Observatory, P.O. Box 296, Epping, N.S.W. 1710, Australia*

²*Research School of Astronomy and Astrophysics, The Australian National University, Private Bag, Weston Creek, A.C.T. 2611, Australia*

³*Astronomy, MS 105-24, Robinson Lab, Caltech, Pasadena, CA 91125, USA*

⁴*Universities Space Research Association*

⁵*NASA Marshall Space Flight Center, ES-84, Huntsville, AL 35812, USA*

⁶*Royal Observatory, Blackford Hill, Edinburgh EH9 3HJ, UK*

⁷*School of Physics A29, University of Sydney, N.S.W. 2006, Australia*

⁸*Astronomical Institute “Anton Pannekoek”, University of Amsterdam, & Center for High Energy Astrophysics, Kruislaan 403, 1098 SJ Amsterdam, The Netherlands*

⁹*Physics Department, University of Alabama in Huntsville, Huntsville, AL 35899, USA*

22 February 2000

ABSTRACT

SN 1998bw holds the record for the most energetic Type Ic explosion, one of the brightest radio supernovae and probably the first supernova associated with a γ -ray burst. In this paper we present spectral observations of SN 1998bw observed in a cooperative monitoring campaign using the AAT, UKST and the SSO 2.3-m telescope. We investigate the evolution of the spectrum between 7 and 94 days after V-band maximum in comparison to well-studied examples of Type Ic SNe in order to quantify the unusual properties of this supernova event. Though the early spectra differ greatly from observations of classical Ic SNe, we find that the evolution from the photospheric to nebular phases is slow but otherwise typical. The spectra differ predominantly in the extensive line blending and blanketing which has been attributed to the high velocity of the ejecta. We find that by day 19, the absorption line minima blueshifts are 10% – 50% higher than other SNe and on day 94 emission lines are 45% broader, as expected if the progenitor had a massive envelope. However, it is difficult to explain the extent of line blanketing entirely by line broadening, and we argue that additional contribution from other species is present, indicating unusual relative abundances or physical conditions in the envelope.

Key words: supernovae: general – stars: evolution – supernovae: general – supernovae: individual: SN 1998bw – gamma-rays: bursts

1 INTRODUCTION

1.1 SN/GRB association

The suggestion that type Ic supernova (SN) 1998bw is the optical counterpart of γ -ray burst (GRB) 980425 has forced us to rethink the mechanics of both SNe and GRBs. The association of SN 1998bw and GRB980425 is supported by the extremely low probability of a chance coincidence (Galama et al. 1998) and particularly by the peculiar observational characteristics of SN 1998bw. Patat & Piemonte (1998a) have classified SN 1998bw as a Type Ic supernova, since spectral lines due to helium, silicon and hydrogen are weak or absent in the early spectra. However, at $M_B = -18.88$ at maximum (Galama et al. 1998), this object was three times

brighter than the average SN Ic, and early spectra show extremely broad lines and unusual line ratios (Lidman et al. 1998). SN 1998bw rivals the brightest radio supernovae yet observed and radio observations indicate that the shock of the explosion was relativistic (Kulkarni et al. 1998; Wieringa et al. 1998). Assuming the association and the low redshift (Tinney et al. 1998) the GRB was also unusual – at least 4 orders of magnitude fainter than other GRBs (Galama et al. 1998).

To what extent SNe are associated with GRBs is unclear due to the low numbers of well-observed Ib/c SNe and the large positional error of most GRBs. Wang & Wheeler (1998) argue that all Ib/c could produce GRBs, but due to beaming we would see only a fraction. Kippen et al.

Table 1. Spectral observations of SN 1998bw.

UT Date	Age ¹ (days)	Telescope	Instrument	λ Coverage (Å)	λ Res. (Å FWHM)	Observer
98/05/19.7	7	2.3m	Nasmyth B Imager	3600-10200	14	Jones, Bessell
98/05/20.6	8	UKST	FLAIR	4001-7151	13	Russell, Parker
98/05/23.8	11	2.3m	Nasmyth B Imager	3600-9990	14	Jones, Bessell
98/05/23.8	11	AAT	2dF	3600-7900	12	Stathakis, Lewis
98/05/27.8	15	AAT	2dF	4035-6263	5	Lewis
98/05/29.7	17	UKST	FLAIR	3950-7202	13	Hartley, Parker
98/05/30.7	18	UKST	FLAIR	3950-7202	13	Hartley, Parker
98/05/31.8	19	AAT	RGO	3520-9300	5	Stathakis, James
98/06/16.6	35	UKST	FLAIR	5700-7560	12	Hartley, Parker
98/06/26	45	2.3m	DBS	3850-7590	6	Germany, Schmidt
98/08/14	94	2.3m	DBS	3800-7550	6	Germany, Schmidt

¹Age is given relative to the date of visual maximum, 1998 May 12.2 (Galama et al. 1998).

(1998) found no evidence for an association between SNe and strong GRBs, and Bloom et al. (1998) model the radio signature and suggest that 1% of GRBs are produced by SNe. If SN 1998bw is a member of a previously unobserved subclass of GRB progenitors (Iwamoto et al. 1998), we have a rare opportunity to test and refine our understanding of SNe and GRBs.

1.2 Type Ib/c supernovae

Supernovae of type Ib/c are identified by their early optical spectra, which lack the deep Si II absorption feature seen at 6150 Å in Ia spectra and the prominent hydrogen lines of Type II SNe. Unlike Ia SNe, Ib/c objects are radio emitters and are typically fainter at maximum by $M_B \sim 1.5$ magnitudes (Filippenko 1997). The parent galaxies of these events (Sbc or later) and heterogeneity of this class suggest that Ib/c SNe are powered by the same mechanism as Type II SNe – core collapse of a massive star. The progenitors of Ib/c SNe have been modelled as Wolf-Rayet stars which have lost their hydrogen envelopes either via close binary interaction (Nomoto, Iwamoto & Suzuki 1995), through a strong stellar wind (Woosley, Langer & Weaver 1993) or a combination of the two mechanisms (Woosley, Langer & Weaver 1995).

This class is further divided into Ib SNe with strong helium lines in the early spectra, and Ic where helium lines are weak or absent. Opinion varies as to the relationship between Ic and helium-rich Ib SNe and whether there is a smooth or bimodal variation of He I strengths (Filippenko et al. 1995; Clocchiatti et al. 1996). Ic progenitors may have lost their helium envelopes in another stage of mass loss, leaving a bare CO star at core collapse (Harkness et al. 1987; Nomoto et al. 1995) or the helium envelope may be poorly mixed with the Ni⁵⁶ (Woosley & Eastman 1997). Piemonte (2000) stresses the spectral and photometric variation for both Ib and Ic SNe. These variations indicate a wide range of ejecta mass, which would depend on the main sequence mass of the progenitor and its mass loss history as well as secondary parameters such as metallicity and convection (Woosley et al. 1995). In general, ejecta mass is expected to be small compared to other SN types.




Figure 1. Comparison of *B*-band (top panel) and *R*-band (bottom panel) spectrophotometry from the corrected spectra with published photometry. Both bands show scatter with $\sigma = 6\%$.

1.3 SN 1998bw

The models for SN 1998bw presented to date tend to fall into two classes – an intrinsically energetic event or hypervelocity (Iwamoto et al. 1998; Woosley et al. 1999) with a massive progenitor star and more normal SNe artificially brightened by beaming (Wang & Wheeler 1998). Radio observations suggest that material has been ejected irregularly by a central engine (Li & Chevalier 1999). All models agree in requiring some form of non-symmetric geometry, possibly produced by an asymmetric explosion.

With such diverse interpretations, it is important that all available data are used to provide observational limits for the models. In this paper we present the results of a cooperative spectral monitoring campaign carried out at Siding Spring, Australia, on the AAT, UKST and SSO 2.3-m telescope between May and November 1998 and compare the spectral evolution and velocity shifts of SN 1998bw with other well-observed Ic SNe.

2 OBSERVATIONS

Spectral observations of SN 1998bw were made in Director's override time and service time on the Anglo-Australian Telescope (AAT) and UK Schmidt Telescope (UKST), and with the cooperation of scheduled observers on the Siding Spring Observatories (SSO) 2.3-m telescope. Unfortunately, the site experienced the poorest observing statistics of the decade and $> 60\%$ of allocated time was lost. We obtained useful spectra at 10 epochs which span 7 to 94 days past V_{max} (or 23 to 110 days past the GRB event). Observational details are given in Table 1.

Observations were made using the scheduled instruments for the telescopes which included both conventional long-slit spectrographs (DBS, RGO) and fibre-fed spectrographs (2dF, FLAIR). The Nasmyth B Imager at the 2.3-m was used as a long-slit spectrograph by inserting blue and red grisms and an order-sorting filter. Long-slit data were processed in the usual way using the FIGARO data reduction package (Shortridge et al. 1997). A first order correction has been made for background emission from the galaxy, but narrow emission from underlying H II regions remain in the spectra. At least two wavelength regions were observed on each of the long-slit data epochs, and spectra were combined by normalising to match the overlap regions.

Data taken on FLAIR were processed using the IRAF data reduction package as outlined in Drinkwater & Holman (1996). 2dF observations were extracted using the S-DIST and C-DIST utilities from FIGARO. Sky emission was removed using neighbouring fibres. It was not possible to correct the 2dF or FLAIR data for background galaxy emission, but the level of contamination was low during this period with the exception of the narrow nebular lines.

Telluric absorption lines have been removed from the data using observations of low-metallicity stars observed on days 7, 11 and 19, scaled where necessary to match the strongest O₂ features. Considering the poor conditions it is probably not surprising that telluric line ratios were variable and weak residuals can be seen in the spectra, particularly around 7500 to 8100 Å and at $\lambda > 9500$ Å.

Long-slit observations were corrected for instrumental response using a contemporaneous observation of a spectrophotometric standard. Fibre observations did not include accompanying standards, so a linear interpolation between bracketing long slit observations were used to correct these data. As these observations were made in non-photometric conditions and through narrow fibres or slits (~ 2 arcsec), correction to absolute flux was made by scaling to match the V-band light curve (Galama et al. 1998; McKenzie & Schaefer 1999) (the V-band was not covered on day 35 so the R-band was used). The success of this method was checked by comparing the photometry in the B and R bands with spectrophotometry from the corrected spectra, measured using filter responses from Bessell (private communication). Both B and R bands resulted in a mean ratio of 0.95, with $\sigma = 6\%$ (Figure 1). Errors at the edge of the spectra are likely to be larger and in general line fluxes from this data set should be regarded with caution. For the purpose of comparison, spectra shown in this paper have been normalised by the V-band photometry, and shifted to rest wavelengths using $vz = 2580$ km s⁻¹ as derived from the narrow H II region emission. Both




Figure 2. Spectral evolution of SN 1998bw. Ages are shown relative to date of visual maximum, 1998 May 12. Spectra have been normalised by the V-band photometry (see text) and offset in flux for the purpose of comparison.

the fluxed and normalised spectra are available via anonymous ftp at <ftp.aao.gov.au/pub/local/ras/98bw>.

3 SPECTRAL EVOLUTION

Early spectral evolution of SN 1998bw has been presented in Iwamoto et al. (1998) (days -9 to +11). In Figure 2 the spectral evolution of SN 1998bw is shown between days 7 and 94. During this period, spectra are dominated by a strong continuum peaking around 5400 Å, with a small number of broad features which become increasingly dominant relative to the continuum. Line widths remain approximately stable. Our observation on day 94 agrees qualitatively with the description by Patat & Piemonte (1998b) of the spectrum on day 123.

The breadth of the spectral features and the uncertainty in continuum level hampers detailed analysis. In this paper we present preliminary results by looking at the most notable features of the spectrum in comparison to classic Ic SNe SN 1983V (Clocchiatti et al. 1997), SN 1987M (Filippenko, Porter & Sargent 1990) and SN 1994I (Clocchiatti et al. 1996; Filippenko et al. 1995), and to SN 1997ef which has unusually broad lines and bears the closest resemblance to SN 1998bw (Iwamoto et al. 2000).




Figure 3. Left: SN 1998bw on days 7 to 11 and days 17 to 19. Right: Spectra of other Ic SNe at similar epochs. All spectra have been corrected for the redshift of the parent galaxy. Positions of median line centres shifted by -10000 km s^{-1} relative to rest are shown in both panels for the Fe II multiplet $\lambda 4274$, Mg II $\lambda 4481$, Fe II $\lambda\lambda 4923, 5018, 5169$, Na I $\lambda\lambda 5890, 5896$ and Si II $\lambda\lambda 6347, 6371$.




Figure 4. Left: Comparison of spectra taken on days 19, 45 and 94. Spectra have been normalised as in previous plots and shifted to rest at $z = 0.0086$. In addition, nebular lines from nearby H II regions have been removed. Right: Spectra of other Ic SNe at similar epochs. Median rest wavelengths are indicated for Mg I] $\lambda 4571$, Fe II $\lambda 5215$, Na I $\lambda\lambda 5890, 5896$, [O I] $\lambda\lambda 6300, 6363$ and Ca II] $\lambda\lambda 7298, 7323$.

Table 2. Absorption minima measurements.

Line	λ_r (Å)	98bw 19d	83V 13d	87M 11d	94I 6d
Ca II	3950	3750 ± 20^1 -15.2 ± 1.5^2	3760 ± 20 -14.4 ± 1.6		3760 ± 10 -14.4 ± 0.8
Fe II	4274	4130 ± 30 -10 ± 2	4150 ± 20 -8.7 ± 1.6	4060 ± 30 -15 ± 2	4170 ± 30 -7 ± 2
Mg II	4481	4330 ± 30 -10 ± 2	4370 ± 20 -7.4 ± 1.4	4360 ± 20 -8.1 ± 1.4	4380 ± 20 -6.8 ± 1.4
Fe II	4555	4330 ± 30 -15 ± 2	4370 ± 20 -12.2 ± 1.4	4360 ± 20 -12.8 ± 1.4	4380 ± 20 -11.5 ± 1.4
Fe II	4923	4700 ± 20 -13.6 ± 1.3	4780 ± 10 -8.7 ± 0.6	4750 ± 10 -10.5 ± 0.6	4800 ± 50 -8 ± 3
Fe II	5018	4820 ± 30 -12 ± 2	4780 ± 10 -14.2 ± 0.6	4820 ± 10 -11.8 ± 0.6	4800 ± 50 -13 ± 3
Fe II	5169	4960 ± 30 -12 ± 2	5000 ± 20 -9.8 ± 1.2	4980 ± 10 -11.0 ± 0.6	5020 ± 10 -8.6 ± 0.6
Na I	5893	5660 ± 30 -11.9 ± 1.5	5710 ± 10 -9.3 ± 0.5	5677 ± 5 -11.0 ± 0.3	5710 ± 7 -9.3 ± 0.4
Si II	6357	6190 ± 30 -7.9 ± 1.4	6240 ± 20 -5.6 ± 1	6152 ± 5 -9.7 ± 0.3	6280 ± 20 -3.7 ± 1.0
O I	7774			7490 ± 10 -11.0 ± 0.4	7560 ± 20 -11.4 ± 1.1

¹Position of absorption minima in wavelength (Å).²Relative velocity (10^3 km s^{-1}).

3.1 Days 7-19

Eight of our spectra fall within the photospheric period, during which the B-band and V-band light curves were declining steeply prior to the radioactive tail. Spectra show little change over the period in the range 4000 to 7000 Å, as seen in Figure 3 (left panel), where spectra have been binned in wavelength and time to improve the signal-to-noise ratio. The height of the continuum is poorly defined as overlapping P-Cygni profiles result in line blanketing over much of the spectral range – apparent peaks are merely regions of relatively low opacity (Iwamoto et al. 1998). In the right panel of Figure 3 spectra of other supernovae are shown at similar epochs. Qualitatively SN 1998bw is very different from the other supernovae, with merged absorption blueward and redward of 5300 Å. SN 1998bw lacks the spectral detail and the overall effect is that of heavy smoothing. However, detailed comparison between the SNe spectra show that the same bands can be identified, and most differences in SN 1998bw can be attributed to the large line widths.

In Table 2, the wavelengths of the observed minima in SN 1998bw (day 19) are compared with those measured from other SNe, and converted to relative velocities (in units of 10^3 km s^{-1}) for suggested line identifications (see below).

Estimates of measurement errors are shown, reflecting how distinct the minima are for each line. SN 1998bw blueshifts are $\sim 30\%$ higher than SN 1983V and $\sim 50\%$ higher than SN 1994I, but are comparable with SN 1987M.

The absorption band between 4100 – 4300 Å is resolved into two minima in SN 1998bw by days 17-19. The two minima are also seen in SN 1983V and SN 1994I. This absorption band is present but not resolved in SN 1997ef. In SN 1987M the band extends blueward compared to the other SNe. The absorption band is attributed to Fe II blends, centred at rest wavelengths of 4274 Å and 4555 Å (Clocchiatti et al. 1997). An alternative identification for the second minimum is Mg II $\lambda 4481$ (Filippenko 1997), which gives velocities which are more in agreement with other features for all but SN 1987M. Other species identified in this region are Ti II and C II (Baron et al. 1996) and Cr II (Iwamoto et al. 1998).

Fe II is also the usual identification for the absorption band between 4700 Å and 5200 Å. In SN 1987M the feature is resolved into three minima which match well with Fe II $\lambda\lambda 4923, 5018, 5169$ (multiplet 42). In SN 1983V and SN 1994I only two minima are resolved, identified as 5018 Å and 5169 Å (Clocchiatti et al. 1997; Iwamoto et al. 1998; Iwamoto et al. 2000). However, we find that the identifi-




Figure 5. Top: Difference between spectra of SN 1998bw observed on day 45 and 94, after scaling by the V-band photometry. Narrow lines from nearby H II regions have been removed. Excess emission from broad nebular lines are visible as SN 1998bw enters the supernebular phase. Bottom: Difference between day 36 and 56 spectra of SN 1994I.

cation of the bluer line as 4923 Å results in better agreement with other features. In SN 1998bw the band is only marginally resolved on day 19, with 4923 Å unusually dominant. In SN 1997ef the line ratios are more typical, and the band is resolved into two minima (Iwamoto et al. 2000).

In Ic SNe spectra, Na I $\lambda\lambda$ 5890, 5896 absorption typically strengthens later than the Fe II features, becoming dominant around day 30 and fading by day 100. The line emerges more slowly in SN 1998bw compared to SN 1983V, SN 1987M and SN 1994I, but is comparable to SN 1997ef. The Na I blueshift is similar to Fe II blueshifts for all four SNe in Table 2. The presence of He I λ 5876 discussed for other Ic SNe (Clocchiatti et al. 1996; Clocchiatti et al. 1997) cannot easily be ascertained for SN 1998bw since the weaker line would be severely blended with Na I.

The Si II $\lambda\lambda$ 6347, 6371 absorption shows the largest shift in velocity during this period, with minima velocities of -11600 km s $^{-1}$ on day 11 and -7900 km s $^{-1}$ on day 19. In Table 2, the Si II line has the lowest blueshift of the measured lines for all SNe. The line profile of the Si II feature is similar to that of SN 1983V, suggesting that there may be contribution from a second line resolved in SN 1983V, SN 1994I and SN 1997ef and identified by Clocchiatti et al. (1997) as O I λ 6158.

3.2 Days 19-94

Late-time spectral evolution of SN 1998bw is shown in Figure 4 (left panel). During this period the light curve

Table 3. Velocity shifts and widths during the nebular phase.

Line	λ_r	98bw (94d)	94I (56d)	87M (62d)	87M (96d)
Mg I]	4571	-1.4^1 11.9 ²	+0.9 8.4	+0.2 8.6	-1.0 7.3
[O I]	6315	+0.7 11.4	-1.2 7.8		-1.9 10.0
Ca II]	7308	-4.0 12.1	-1.6 8.2	-2.7 7.5	-2.6 6.5
Na I]	5893 ³	-10.1 11.1	-11.3 9.0	-10.0 5.5	-9.9 4.5

¹Velocity shift of line centre (10³ km s $^{-1}$).

²Width of features at half height (10³ km s $^{-1}$).

³Absorption component.

is decaying linearly via radioactive decay (McKenzie & Schaefer 1999), and as expected we see a transition from an absorption-dominated photospheric spectrum to an emission-dominated nebular spectrum, though on day 94 the transition is still not complete. Early nebular spectra of SN 1994I and SN 1987M are shown in comparison to SN 1998bw in Figure 4 (right panel). SN 1998bw evolves slowly compared to the other two SNe – the day 94 spectrum bears a closer resemblance to SN 1987M on day 62 than day 96, and is very similar to SN 1994I on day 56. In order to separate the new emission from the other features in SN 1998bw, the day 45 spectrum has been subtracted from the day 94 spectrum, after scaling by the V-band photometry. Likewise, day 36 has been subtracted from day 56 for SN 1994I. The results are compared in Figure 5.

Velocity shifts of the main features are given in Table 3. [O I] $\lambda\lambda$ 6300, 6363 is blueshifted in SN 1987M and SN 1994I. In SN 1998bw on day 94 the line profile is symmetric, but in the residual emission it is also blueshifted by ~ -500 km s $^{-1}$. Ca II] $\lambda\lambda$ 7291, 7323 has a similar blueshift in SN 1987M and SN 1994I. In SN 1998bw the blueshift is significantly greater, but in the residual spectrum the profile is similar to SN 1994I, so apparent differences are probably due to contribution from the persisting photospheric features. The Ca II] to [O I] ratio is greater in SN 1987M than in SN 1994I and SN 1998bw, which has been attributed to a difference in the relative abundances of calcium and oxygen (Filippenko et al. 1995). While Mg I] λ 4571 is the usual identification for the emission peak at 4500 Å (Filippenko 1997), an alternative identification is given by Patat et al. (1998b) as Fe II λ 4555, and both transitions give an adequate fit to the line with a symmetric profile. We adopt the Mg I] identification for this paper.

Approximate line widths have been measured using ABLINE in FIGARO (Table 3). Line widths are similar for the Mg I], [O I] and Ca II] emission for each supernova. The mean width of these lines in SN 1998bw on day 94 is 11600 ± 400 km s $^{-1}$ which is $\sim 45\%$ broader than SN 1987M and SN 1994I. Na I emerges as a P-Cygni profile between days 19 and 94, evolving more slowly in equivalent width and in emission to absorption ratio than SN 1987M and




Figure 6. The 7000 to 9000 Å region of Ic SNe spectra is dominated by the O I $\lambda 7774$ multiplet and Ca II $\lambda\lambda 8498, 8542, 8662$ (rest wavelengths are shown). These lines are blended in SN 1998bw on day 11 (a) and day 19 (b), but are well separated in SN 1987M on day 7 (c). Note that weak residual telluric lines are present in the SN 1998bw spectra, especially around 7600 Å.

SN 1994I. A noticeable difference between SN 1998bw and SN 1987M is the width of the absorption component of Na I, which is only half the width in SN 1987M, and considerably narrower than the emission lines.

The peak at 5200 Å has been tentatively identified as Fe II $\lambda 5215$ (Patat & Piemonte 1998b). This transition is typically seen in older Ia SNe, but not in Ic SNe, and Patat and Piemonte note that its presence would indicate that SN 1998bw was a type Iac SN. However, the feature is also present in SN 1987M and more weakly in SN 1994I during the early nebular phase. It fades relative to Mg I], [O I] and Ca II] at later times. It therefore seems reasonable to assume that the presence of Fe II emission in Ic SNe spectra is typical during the transition from the photospheric to the nebular phase. In the residual spectrum (Figure 5) the feature is weaker relative to the 4500 Å peak and has a markedly different profile. This supports the identification of the 4500 and 5200 Å peaks as transitions of different species, and indicates that the 5200 Å peak is fading and/or is artificially enhanced by a relatively high continuum level.

3.3 7000 – 9000 Å

The red spectral region of SN 1998bw is shown in Figure 6 for days 7 and 19. It is this region which differs most markedly from typical Ic SNe such as SN 1987M (shown on day 7). In these SNe, strong absorption from O I $\lambda 7774$ and Ca II $\lambda\lambda 8498, 8542, 8662$ are well separated by an absorption

free region missing in SN 1998bw. Iwamoto et al (1998) have successfully modelled this region for day –9 with a photospheric velocity of 28000 km s^{–1}. However, by day –1 their model predicts that the two features should be resolved, and that by day 7 O I and Ca II are unblended. Similar evolution is predicted by the direct analysis models of Branch (2000). In SN 1997ef this behaviour is seen, with blended absorption on day 3 and well separated features on day 30 (Iwamoto et al. 2000). In SN 1998bw, however, there is no sign of significant separation as late as day 19, our last epoch covering this region.

While it is presumably possible to fit this region by increasing the mass of the progenitor, and therefore the line widths of the O I and Ca II profiles, there are two limitations which need to be considered. The first is that we do not expect to see redshifted absorption from a simple expanding envelope. Under this assumption, O I cannot contribute to the absorption band redward of 7774 Å, irrespective of the line width. The second limitation is that we detect a shallow dip at ~8100 Å which we identify as the minimum of the Ca II absorption. This feature aligns well with the Ca II $\lambda\lambda 3933, 3968$ line profile, and the velocity of –15300 km s^{–1} is already higher than blueshifts of other lines at this epoch (Table 3).

In order to reproduce the spectrum on day 19 we would require a highly unusual geometry to produce redshifted absorption from O I, or to produce Ca II absorption with a second minimum at around –24000 km s^{–1}. An alternative explanation is the presence of a third component absorbing at around 7700 – 8000 Å. Adequate fits can be produced with lines of rest wavelength 8000 – 8350 Å, and candidate species include C II, C III and N I. A relatively weak contribution from any of these species would result in the blended spectrum we observe. Modelling is required to further investigate this region, and to determine whether unusual abundances, density or temperature distribution can explain the observations.

4 CONCLUSION

During the period between 7 and 94 days after V-band maximum, we have seen that SN 1998bw resembles other Ic SNe sufficiently to support this classification, but has unusually slow spectral evolution. On day 94 we see the emergence of a nebular spectrum, which retains many of the characteristics of the photospheric period. The late onset of the supernebular phase, compared to SN 1987M (62 d) and SN 1994I (56 d), is consistent with the ejection of an unusually large mass, as predicted by lightcurve models (Iwamoto et al. 1998; Woosley et al. 1999).

By day 19, SN 1998bw blueshifts are up to 50% larger than other Ic SNe. However, increased blueshifts alone seem insufficient to explain the unusually smooth and blended spectrum which persists to late times – for instance SN 1998bw on day 19 has similar blueshifts to SN 1987M on day 11, but SN 1987M has well defined spectral features. Emission line widths on day 94 are 45% broader than SN 1994I and SN 1987M and Na I absorption is far broader in SN 1998bw on day 94 than in SN 1987M at similar epochs. The line profiles of SN 1998bw may have disproportionately strong absorption wings – we lack an example of an un-

blended feature at earlier times for confirmation. Using the standard model for a homologously expanding envelope, absorption lines which are broader but of similar blueshift imply that the absorption region in SN 1998bw spans a larger range of velocity space, at both higher and lower velocities than other Ic SNe, as expected for a massive envelope. A closer inspection of the 7000 – 9000 Å region of SN 1998bw suggests that we are also seeing contribution from enhanced line species. Unusually strong lines from species such as N I, C II, C III, Ti II and Cr II may help to produce the extensive line blending. If so, this could indicate an overabundance of these elements, or unusual physical properties of the ejecta.

More work is required in establishing line identifications and spectral characteristics, best done using spectral models. Whether or not SN 1998bw was associated with GRB 980425, it is of great interest as an extreme example of Ic SNe. SN 1997ef bears some resemblance to SN 1998bw and may be an intermediate object between SN 1998bw and classical SNe, though it is too early to say whether we are seeing a bimodal or continuous variation in properties. Thanks to the interest inspired by the γ -ray burst, SN 1998bw has been observed extensively and successful modelling of this object is likely to enhance our understanding of this relatively poorly observed class of supernova.

Acknowledgements

We thank the observers at the AAT, UKST and MSO 2.3-m telescope who kindly donated their time to our project. We thank Alejandro Clocchiatti, Craig Wheeler, Jodie Martin and Paolo Mazzali for providing digital supernova data to compare with our observations, and Peter Meikle, David Branch and Brian Schmidt for helpful discussions.

REFERENCES

- Baron E., Hauschildt P. H., Branch D., Kirshner R. P., Filippenko A. V., 1996, MNRAS, 279, 799, astro-ph/9510070
 Bloom J. S., Kulkarni S. R., Harrison F., Prince T., Phinney E. S., 1998, ApJ, 506, L105, astro-ph/9807050
 Branch D., 2000, in Livio M., ed, Supernovae and Gamma-Ray Bursts, in press, astro-ph/9906168
 Clocchiatti A., Wheeler J. C., Brotherton M. S., Cochran A. L., Wills D., Barker E. S., Turatto M., 1996, ApJ, 462, 462
 Clocchiatti A. et al., 1997, ApJ, 483, 675
 Drinkwater M., Holman B., 1996, FLAIR Data Reduction with IRAF, http://www.aao.gov.au/local/www/cgt/flair_iraf/flair_iraf.html
 Filippenko A. V., 1997, ARA&A, 35, 309
 Filippenko A. V. et al., 1995, ApJ, 450, L11
 Filippenko A. V., Porter A. C., Sargent W. L. W., 1990, AJ, 100, 1575
 Galama T. J. et al., 1998, Nature, 395, 670, astro-ph/9806175
 Harkness R. P. et al., 1987, ApJ, 317, 355
 Iwamoto K. et al., 1998, Nature, 395, 672, astro-ph/9806382
 Iwamoto K., Nakamura T., Nomoto K., Mazzali P. A., Danziger I. J., Garnavich P., Kirshner R., Jha S., Balam D., Thorstensen J., 2000, ApJ, in press, astro-ph/9807060
 Kippen R. M. et al., 1998, ApJ, 506, L27, astro-ph/9806364
 Kulkarni S. R. et al., 1998, Nature, 395, 663, astro-ph/9807001
 Li Z.-Y., Chevalier R. A., 1999, ApJ, 526, 716, astro-ph/9903483
 Lidman C., Doublier V., Gonzalez J. -F., Augusteijn T., Harnaut O. R., Boehnhardt H., Patat F., Leibundgut B., 1998, IAU 6895
 McKenzie E. H., Schaefer B. E., 1999, PASP, 111, 964, astro-ph/9904397
 Nomoto K., Iwamoto K., Suzuki T., 1995, Physics Reports 256, 173
 Patat F., Piemonte A., 1998a, IAU 6918
 Patat F., Piemonte A., 1998b, IAU 7017
 Piemonte A., 2000, in Phillips M., ed, SN 1987A: Ten Years Later, in press
 Shortridge K., Meyerdierks H., Currie M., Clayton M., 1997, PPARC Starlink User Note, 86.13
 Tinney C., Stathakis R., Cannon R., Galama T., 1998, IAU 6896
 Wang L., Wheeler J. C., 1998, ApJ, 504, L87, astro-ph/9806212
 Wieringa M., Frail D. A., Kulkarni S. R., Higdon J. L., Wark R., Bloom J. S., BeppoSAX GRB Team, 1998, IAU 6896
 Woosley S. E., Eastman R. G., 1997, in Ruiz-Lapuente R., Canal R., Isern I., eds, Thermonuclear Supernovae, NATA ASI series, 821
 Woosley S. E., Eastman R. G., Schmidt B. P., 1999, ApJ, 516, 788, astro-ph/9806299
 Woosley S. E., Langer N., Weaver T. A., 1993, ApJ, 411, 823
 Woosley S. E., Langer N., Weaver T. A., 1995, ApJ, 448, 315